

# Surgery Obstructions from Khovanov homology

Liam Watson  
Université du Québec à Montréal


`www.cirget.uqam.ca/~liam`

Georgia Topology Conference  
May 21, 2009

# Involutions and tangles

Let  $K \hookrightarrow S^3$  be a strongly invertible knot.

Then there is an involution  $f$  on the knot complement  $M = S^3 \setminus \nu(K)$  with one dimensional fixed point set (a pair of arcs) meeting the boundary transversely in 4 distinct points.



Note that the quotient  $M/f$  is homeomorphic to a 3-ball.



## Definition

For a strongly invertible knot  $K \hookrightarrow S^3$ , the *associated quotient tangle* is the pair  $T = (B^3, \tau)$ , where  $\tau$  is the image of the fixed point set of  $f$  in the quotient  $M/f \cong B^3$ .


As a result the knot complement is a two-fold branched cover:

$$M \cong \Sigma(B^3, \tau).$$

# Example: the figure eight




# Example: the figure eight




## Example: the figure eight

Tangles, in this setting, are considered up to homeomorphism of the pair  $(B^3, \tau)$ :



In particular, such homeomorphisms need not fix the boundary.


# The preferred representative



By construction, the *denominator* closure of  $T$  – denoted  $\tau(\frac{1}{0})$  – corresponds to the trivial surgery on  $K$  (notice that  $\tau(\frac{1}{0})$  is the trivial knot).



There is a preferred choice of representative for the associated quotient tangle so that the *numerator* closure of  $T$  – denoted  $\tau(0)$  – corresponds to the zero surgery in the cover:  $S_0^3(K) \cong \Sigma(S^3, \tau(0))$ .



In particular, with this notation  $\tau(n)$  is obtained by adding  $n$  half-twists so that  $S_n^3(K) \cong \Sigma(S^3, \tau(n))$ .


# Montesinos' trick

In general,

$$S^3_{p/q}(K) \cong \Sigma(S^3, \tau(\frac{p}{q}))$$

where the link  $\tau(\frac{p}{q})$  is obtained by attaching a rational tangle. For example:

$$\tau(\frac{13}{10}) = \tau[1, 3, 3] =$$



With the observation that

$$S_{p/q}^3(K) \cong \Sigma(S^3, \tau(\frac{p}{q}))$$

in hand, the idea is to apply the Khovanov homology of  $\tau(\frac{p}{q})$  as an obstruction to exceptional Dehn surgeries on  $K$ .

The reduced Khovanov homology is a relatively  $\mathbb{Z} \oplus \mathbb{Z}$ -graded group  $\widetilde{\text{Kh}}(L)$  associated to a link  $L \hookrightarrow S^3$ . We work over  $\mathbb{F} \cong \mathbb{Z}/2\mathbb{Z}$ , with primary (cohomological) grading  $\delta$  and secondary (Jones, quantum) grading  $q$ . These grading conventions are non-standard:



## Theorem (Khovanov)

Let  $u = \delta + q$ , then there exists an absolute  $\mathbb{Z} \oplus \frac{1}{2}\mathbb{Z}$ -grading on  $\widetilde{\text{Kh}}(L)$  so that


$$V_L(t) = \sum_{u,q} (-1)^u t^q \text{rk } \widetilde{\text{Kh}}_q^u(L)$$

where  $V_L(t) \in \mathbb{Z}[t^{\frac{1}{2}}, t^{-\frac{1}{2}}]$  is the Jones polynomial.

# Example: the trefoil



$$\delta = u - q \longrightarrow$$



## Definition

The homological width of a link  $L$  is given by the number of  $\delta$ -gradings supporting  $\widetilde{\text{Kh}}(L)$ . That is if

$$\bigoplus_{\delta} \widetilde{\text{Kh}}^{\delta}(L) \cong \mathbb{F}^{b_1} \oplus \dots \oplus \mathbb{F}^{b_k},$$

for  $b_{\delta} \geq 0$  and  $b_1, b_k > 0$ , write  $w(L) = k$ .


Notice that

$$\left| \sum_{\delta} (-1)^{\delta} \text{rk } \widetilde{\text{Kh}}^{\delta}(L) \right| = |H_1(\Sigma(S^3, L); \mathbb{Z})|$$

since  $|V_L(-1)| = \det(L) = |H_1(\Sigma(S^3, L); \mathbb{Z})|$ .


# Examples

1
1
1



$$w(3_1) = 1$$

	1
	1
1	1
1	1
1	



$$w(10_{124}) = 2$$

## Theorem 1 (W.)

*If  $\Sigma(S^3, L)$  has finite fundamental group then  $w(L) \leq 2$ .*

As a first step, compare:

## Theorem

*If  $\Sigma(S^3, L)$  is a lens space then  $w(L) = 1$ .*

## Proof.

Hodgson and Rubinstein show that if  $\Sigma(S^3, L)$  is a lens space then  $L$  is a non-split two-bridge link; Lee proved that non-split alternating links – in particular two-bridge links – are thin. □

# Manifolds with finite fundamental group

- By the orbifold theorem, having a finite fundamental group is equivalent to admitting elliptic geometry in this setting (Thurston, see Boileau-Porti).
- Manifolds with elliptic geometry are all Seifert fibered: they are either lens spaces (see previous theorem) or have base orbifold  $S^2(2, 2, n)$  for  $n > 1$  or  $S^2(2, 3, n)$  for  $n = 3, 4, 5$  (Seifert, see Scott).
- These manifolds may be constructed by considering Dehn fillings of the twisted  $I$ -bundle over the Klein bottle (base  $D^2(2, 2)$ ) or the trefoil complement (base  $D^2(2, 3)$ ) (Heil, Montesinos).
- This construction is such that the branch set in each case is recovered, and this branch set is unique (Montesinos, Boileau-Otal).

# Manifolds with finite fundamental group

In summary, there exists a set of links  $\mathcal{L}$  for which  $L \in \mathcal{L}$  if and only if  $\pi_1(\Sigma(S^3, L))$  is finite.

To prove Theorem 1, we need to see that this collection of branch sets has relatively tame Khovanov homology, in the sense that  $w(L) \leq 2$  whenever  $L \in \mathcal{L}$ .

This will rely on a particular form of **stability** enjoyed by Khovanov homology.

Let  $K \hookrightarrow S^3$  be a strongly invertible knot so that  $S^3_{p/q}(K) \cong \Sigma(S^3, \tau(\frac{p}{q}))$ . Define

$$w_K = \min_{\frac{p}{q} \in \mathbb{Q}} \{w(\tau(\frac{p}{q}))\}.$$

## Theorem 2 (W.)

*If  $w_K > 1$  then  $K$  does not admit lens space surgeries, and if  $w_K > 2$  then  $K$  does not admit finite fillings. Moreover, if  $T$  is **generic** then  $w_K$  is determined on a finite collection of integer fillings by **stability**.*

# The skein exact sequence

$$\begin{array}{ccc}
 & \widetilde{\text{Kh}}(\text{X}) & \\
 \nearrow & & \searrow \\
 \widetilde{\text{Kh}}(\text{O})[-\frac{c}{2}, \frac{3c+2}{2}] & \xleftarrow{[1,0]} & \widetilde{\text{Kh}}(\text{O})[-\frac{1}{2}, \frac{1}{2}]
 \end{array}$$


Where  $c = n_-(\text{O}) - n_-(\text{X})$  and  $\widetilde{\text{Kh}}_q^\delta(L)[i, j] = \widetilde{\text{Kh}}_{q-j}^{\delta-i}(L)$ .

Or, as a mapping cone:

$$\widetilde{\text{Kh}}(\text{X}) \cong H_* \left( \widetilde{\text{Kh}}(\text{O})[-\frac{1}{2}, \frac{1}{2}] \rightarrow \widetilde{\text{Kh}}(\text{O})[-\frac{c}{2}, \frac{3c+2}{2}] \right)$$

# A mapping cone for integer surgeries

Now when applying this to the link  $\tau(m+1)$  we have:



# A mapping cone for integer surgeries

So that

$$\widetilde{\text{Kh}}(\tau(m+1)) \cong H_* \left( \widetilde{\text{Kh}}(\tau(m))[-\frac{1}{2}, \frac{1}{2}] \rightarrow \widetilde{\text{Kh}}(\tau(\frac{1}{0}))[-\frac{c}{2}, \frac{3c+2}{2}] \right)$$

where  $\tau(\frac{1}{0})$  is the trivial knot and  $c = c_\tau + m$  with

$$c_\tau = n_- \left( \left( \text{link}(T) \right) \right) - n_- \left( \text{link}(T) \right)$$

$$\begin{aligned} & \widetilde{\text{Kh}}(\tau(m+1)) \\ & \cong H_* \left( \widetilde{\text{Kh}}(\tau(m))[-\frac{1}{2}, \frac{1}{2}] \rightarrow \mathbb{F}[-\frac{c_\tau}{2}, \frac{3c_\tau+2}{2}][0, m][-\frac{m}{2}, \frac{m}{2}] \right) \end{aligned}$$

# A mapping cone for integer surgeries

## Stability Lemma

For any integer  $m$ , and positive integer  $n$ ,

$$\widetilde{\text{Kh}}(\tau(m+n)) \cong H_* \left( \widetilde{\text{Kh}}(\tau(m)) \rightarrow \bigoplus_n \widetilde{\text{Kh}}(\tau(\frac{1}{0})) \right)$$

as a relatively  $\mathbb{Z} \oplus \mathbb{Z}$ -graded group. More precisely, there exist explicit constants  $x$  and  $y$  and an identification


$$\bigoplus_{q=0}^{n-1} \widetilde{\text{Kh}}(\tau(\frac{1}{0}))[x, y][0, q] \cong \mathbb{F}[\mathbb{Z}/n\mathbb{Z}]$$

as graded  $\mathbb{F}$ -vector spaces so that

$$\widetilde{\text{Kh}}(\tau(m+n)) \cong H_* \left( \widetilde{\text{Kh}}(\tau(m)) \rightarrow \mathbb{F}[\mathbb{Z}/n\mathbb{Z}] \right).$$


# Example: the figure eight revisited

		1
		1
	1	1
	1	1
1	1	
	1	
	1	




# Example: the figure eight revisited

		1
		1
	1	1
	1	1
	1	
	1	
	1	
	1	




# Example: the figure eight revisited

		1
		1
	1	1
	2	1
	1	
	1	
	1	




# Example: the figure eight revisited

		1
		1
	1	1
	1	
	2	1
	1	
	1	
	1	




# Example: the figure eight revisited

		1
		1
	2	1
	1	
	2	1
	1	
	1	
	1	



# Example: the figure eight revisited

		1
	1	1
	2	1
	1	
	2	1
	1	
	1	
	1	



## Example: the figure eight revisited

Notice that  $w(\tau(n)) = 2$  for  $n > 0$ , and  $w(\tau(n)) = 3$  for  $n \leq 0$  as a consequence of the stability lemma.

By the cyclic surgery theorem, a lens space surgery on  $S^3$  arises as an integer surgery.

Therefore, we recover the well known fact that the figure eight does not admit lens space surgeries:

$$\widetilde{\text{Kh}}(\tau(0)) \cong \mathbb{F} \oplus \mathbb{F}^5 \oplus \mathbb{F}^4$$

implies that  $w > 1$  for branch sets associated to integer surgeries.

## Lemma

For  $N \gg 0$  the exact sequence for  $\widetilde{\text{Kh}}(\tau(N+1))$  splits so that, ignoring gradings,

$$\widetilde{\text{Kh}}(\tau(N+1)) \cong \widetilde{\text{Kh}}(\tau(N)) \oplus \mathbb{F}.$$

## Lemma

Up to overall shift the generators  $\widetilde{\text{Kh}}(\tau(\frac{1}{0})) \cong \mathbb{F}$ , when they survive in homology, are all supported in a single relative  $\delta$ -grading.

## Definition

For a given strongly invertible knot and preferred associated quotient tangle, define

$$w_{\max} = \max_{n \in \mathbb{Z}} \{w(\tau(n))\}$$

and

$$w_{\min} = \min_{n \in \mathbb{Z}} \{w(\tau(n))\}.$$

## Lemma

*Either  $w_{\max} = w_{\min}$  or  $w_{\max} = w_{\min} + 1$ .*

# An upper bound for width

With a view to proving Theorem 1:

## Proposition

*Let  $K$  be a strongly invertible knot with preferred associated quotient tangle  $T$ . Then  $w(\tau(\frac{p}{q})) \leq w_{\max}$ .*

To prove the proposition, it is natural to introduce

$$\widetilde{\text{Kh}}_{\sigma}(L) \cong \widetilde{\text{Kh}}(L)[- \frac{\sigma(L)}{2}]$$

as an *absolutely*  $\mathbb{Z}$ -graded object where  $\sigma(L)$  is the signature.

**Theorem (Manolescu-Ozsváth)**


$$\widetilde{\text{Kh}}_{\sigma}(\text{X}) = H_* \left( \widetilde{\text{Kh}}_{\sigma}(\text{Y}) \rightarrow \widetilde{\text{Kh}}_{\sigma}(\text{Z}) \right)$$

if  $\det(\text{Y}), \det(\text{Z}) > 0$  and  $\det(\text{X}) = \det(\text{Y}) + \det(\text{Z})$ .

It is possible to prove a variant of this statement when the determinant of one of the resolutions vanishes.

# Resolutions and continued fractions

$$\frac{p}{q} = \frac{13}{10} = [1, 3, 3]$$



$$\frac{p_1}{q_1} = \frac{4}{3} = [1, 3]$$

$$\frac{p_0}{q_0} = \frac{9}{7} = [1, 3, 2]$$

$$\frac{13}{10} = \frac{4+9}{3+7}$$

# Resolutions and continued fractions

In general,

$$\frac{p}{q} = \frac{p_0 + p_1}{q_0 + q_1}$$

when

$$\frac{p}{q} = [a_1, \dots, a_{r-1}, a_r - 1, 1] = [a_1, \dots, a_{r-1}, a_r]$$

and  $\frac{p_0}{q_0}, \frac{p_1}{q_1}$  are the continued fractions

$$[a_1, \dots, a_{r-1}], [a_1, \dots, a_{r-1}, a_r - 1].$$

Since  $\det(\tau(\frac{p}{q})) = |H_1(\Sigma(S^3, \tau(\frac{p}{q})); \mathbb{Z})| = |H_1(S^3_{p/q}(K); \mathbb{Z})| = p$   
we have that

$$\det(\tau(\frac{p}{q})) = \det(\tau(\frac{p_0}{q_0})) + \det(\tau(\frac{p_1}{q_1}))$$

and Manolescu and Ozsváth's theorem may be applied.

# Resolutions and continued fractions

As a result, it is possible to induct in the length  $r$  of the continued fraction to prove that  $w_{\max}$  is an upper bound for  $w(\tau(\frac{p}{q}))$ .

In particular, by successively resolving the final crossing of  $\tau(\frac{p}{q})$  it can be shown that

$$\begin{aligned}w(\tau(\frac{p}{q})) &\leq \max\{w(\tau\lfloor\frac{p}{q}\rfloor), w(\tau\lceil\frac{p}{q}\rceil)\} \\ &= \max\{w(\tau(a_1)), w(\tau(a_1 + 1))\}.\end{aligned}$$

where  $\frac{p}{q} = [a_1, \dots, a_{r-1}, a_r]$ .

# On Quasi-alternating links

## Definition

The set of quasi-alternating links  $\mathcal{Q}$  is the smallest set of such that:

- The trivial knot is an element of  $\mathcal{Q}$ , and
- if  $L$  admits a projection with distinguished crossing  $\times$  for which each resolution gives an element of  $\mathcal{Q}$ , and

$$\det(\times) = \det(\smile) + \det(\frown),$$

then  $L \in \mathcal{Q}$  as well.

## Theorem (Manolescu-Ozsváth)

*Quasi-alternating links are homologically thin.*

## Proposition

Suppose  $S_{p/q}^3(K) \cong \Sigma(S^3, \tau(\frac{p}{q}))$  and  $\tau(N)$  is quasi-alternating for some  $N > 0$ . Then  $\tau(\frac{p}{q})$  is quasi-alternating for all  $\frac{p}{q} \geq N$ .

## Corollary

For large surgery on the trefoil,  $\tau(\frac{p}{q})$  is quasi-alternating. In particular,  $w(\tau(\frac{p}{q})) = 1$  for  $\frac{p}{q} \geq 5$ .

# $w \leq 2$ for manifolds with finite fundamental group

Since  $w_{\max} = w_{\min} + 1 = 2$  for the tangle associated to the trefoil,  $w(L) \leq 2$  for  $\Sigma(S^3, L)$  Seifert fibered with base orbifold  $S^2(2, 3, n)$ .

A similar argument holds for branch sets associated to fillings of the twisted  $I$ -bundle over the Klein bottle to obtain the  $S^2(2, 2, n)$  family.

This proves Theorem 1.

# A lower bound for width

The proof of Theorem 2 depends on similar arguments to establish  $w_{\min}$  as a lower bound for  $w(\tau(\frac{p}{q}))$ .

## Proposition

*Let  $K$  be a strongly invertible knot with **generic** preferred associated quotient tangle  $T$ . Then  $w(\tau(\frac{p}{q})) \geq w_{\min}$ .*

In particular:

$$w_K = w_{\min}$$

A tangle  $T$  is generic if either

- $w_{\max} = w_{\min}$ , OR
- if  $b_k > 1$  where

$$\widetilde{\text{Kh}}(\tau(\ell)) \cong \mathbb{F}^{b_1} \oplus \dots \oplus \mathbb{F}^{b_k}$$

and

$$\widetilde{\text{Kh}}(\tau(\ell + 1)) \cong \mathbb{F}^{b_1} \oplus \dots \oplus \mathbb{F}^{b_k} \oplus \mathbb{F},$$

or

- if  $b_1 > 1$  where

$$\widetilde{\text{Kh}}(\tau(\ell)) \cong \mathbb{F} \oplus \mathbb{F}^{b_1} \oplus \dots \oplus \mathbb{F}^{b_k}$$

and

$$\widetilde{\text{Kh}}(\tau(\ell + 1)) \cong \mathbb{F}^{b_1} \oplus \dots \oplus \mathbb{F}^{b_k}.$$

For example, for the figure eight we had that

$$\widetilde{\text{Kh}}(\tau(0)) \cong \mathbb{F} \oplus \mathbb{F}^5 \oplus \mathbb{F}^4$$

and

$$\widetilde{\text{Kh}}(\tau(+1)) \cong \mathbb{F}^5 \oplus \mathbb{F}^4$$

so that the width *decays* but  $b_1 = 5$  so the tangle is generic.  
Since the figure eight is amphicheiral, we recover:

## Theorem (Thurston)

*The figure eight does not admit finite fillings.*

# A lower bound for width

Suppose that  $w_{\min} = w_{\max} = w$ . Then as before

$$\widetilde{\text{Kh}}_{\sigma}(\tau(\frac{p}{q})) \cong H_* \left( \widetilde{\text{Kh}}_{\sigma}(\tau(\frac{p_0}{q_0})) \rightarrow \widetilde{\text{Kh}}_{\sigma}(\tau(\frac{p_1}{q_1})) \right).$$

Recall that the connecting homomorphism raises  $\delta$ -grading by one:

$$\widetilde{\text{Kh}}(\tau(\frac{p}{q})) \cong H_* \left( \begin{array}{cccc} \mathbb{F}^{b_1} & \mathbb{F}^{b_2} & \cdots & \mathbb{F}^{b_w} \\ & \searrow & \searrow & \searrow \\ \mathbb{F}^{b'_1} & \mathbb{F}^{b'_2} & \cdots & \mathbb{F}^{b'_w} \end{array} \right)$$

By induction in the length of the continued fraction for  $\frac{p}{q}$ ,  
 $w(\tau(\frac{p}{q})) = w$ .

# Determining widths


Notice that  $w(\tau(-)) : \mathbb{Q} \rightarrow \mathbb{N}$  is constant when  $w_{\min} = w_{\max}$ .

After a slightly modified argument when  $w_{\max} = w_{\min} + 1$ , we have  $w(\tau(-)) : \mathbb{Q} \rightarrow \mathbb{N}$  takes values  $\{w_{\min}, w_{\max}\}$  in the generic setting.

This proves Theorem 2: for generic tangles, the minimum width is determined on the integer fillings. That is,

$$w_K = w_{\min}.$$

# Example: the knot $14_{11893}^n$



## Theorem (Ozsváth-Szabó)

If  $K \hookrightarrow S^3$  admits an  $L$ -space surgery then

$$\Delta_K(t) = (-1)^k + \sum_{j=1}^k (-1)^{k-j} (t^{-n_j} + t^{n_j})$$


for  $0 < n_1 < n_2 < \dots < n_k$ .

For example,  $\Delta_{4_1}(t) = -t^{-1} + 3 - t$ .


On the other hand,

$$\Delta_{14_{11893}^n}(t) = t^{-3} - t^{-2} + t^{-1} - 1 + t - t^2 + t^3.$$

Example: the knot  $14_{11893}^n$



			1
			2
			2
		2	3
		4	3
		5	2
	2	7	2
	4	7	1
	5	6	
1	6	5	
2	7	3	
3	5		
3	4		
4	3		
3			
2			
2			



For any  $n$

$$w(\tau(n)) \geq 4$$

so that


$$w_k = 4$$

**Theorem**

$14_{11893}^n$  does not admit finite fillings.


			1
			2
			2
		2	3
		4	3
		5	2
	2	7	2
	4	7	1
	5	6	
1	6	5	
2	7	3	
3	5		
3	4		
4	3		
3			
2			
2			

$$\det(\tau(m+2)) = 7$$




			1
			2
			2
		2	3
		4	3
		5	2
	2	7	2
	4	7	1
	5	6	
1	6	5	
2	7	④	
3	5	①	
3	4		
4	3		
3			
2			
2			

$$\det(\tau(m)) = 9$$



			1
			2
			2
		2	3
		4	3
		5	2
	2	7	2
	4	7	1
	5	6	
1	6	5	
2	7	3	
3	5		
3	4		
4	3		
3			
2			
2			

$$\widetilde{\text{Kh}}(\tau(-7)) \cong \mathbb{F}^{20} \oplus \mathbb{F}^{36} \oplus \mathbb{F}^{39} \oplus \mathbb{F}^{16}$$



			1
			2
			2
		2	3
		4	3
		5	2
	2	7	2
	4	7	1
	5	6	
1	6	5	
2	7	④	
3	5	①	
3	4		
4	3		
3			
2			
2			

$$\widetilde{\text{Kh}}(\tau(-9)) \cong \mathbb{F}^{20} \oplus \mathbb{F}^{36} \oplus \mathbb{F}^{41} \oplus \mathbb{F}^{16}$$

